

Microensing Searches for Extrasolar Planets: Current Status and Future Prospects

B. Scott Gaudi

*Hubble Fellow, School of Natural Sciences, Institute for Advanced Study,
 Einstein Drive, Princeton, NJ, 08540*

Abstract. I review results from, and future prospects for, microlensing searches for extrasolar planets. Analyses of well-sampled microlensing light curves by several collaborations have demonstrated that current searches are sensitive to $\gtrsim 1M_J$ planets with few AU separations from M dwarfs in the Galactic bulge. To date, however, no unambiguous planetary detections have been made. Detailed analysis has shown that this null result implies that $< 33\%$ of typical stars (i.e. M dwarfs) in the Galactic bulge have Jupiter-mass companions with separations between 1.5 and 4 AU, and $< 45\%$ have $3M_J$ companions between 1 and 7 AU. The recent dramatic increase in the number of alerts per year will allow ongoing microlensing searches to probe companion fractions of a few percent within a few years.

1. Introduction

Ultra-precise radial velocity (RV) surveys have revealed over 100 planetary companions to nearby FGKM main-sequence stars.¹ The minimum masses M_p and semi-major axes a of these companions are shown in Figure 3. The number of known planetary companions is now reaching the point where robust statistical inferences about trends within the sample can be made, and thus the study of extrasolar planets is in transition from the discovery phase to the characterization phase.

The ‘classical’ methods of detecting extrasolar planets (RV, astrometry, transits, direct detection) are generally complementary to each other both in the range of semi-major axes a they probe (see Fig. 3), and in the parameters they measure. They also generally suffer from the same set of drawbacks. First, because they rely on light from either the parent star or the planet itself, they are generally limited to nearby systems. Second, they are not currently sensitive to very low-mass planets. For example, the systematic floor of RV surveys is thought to be $\sim 1 \text{ m s}^{-1}$. This implies that Earth, Uranus, and Neptune analogs are probably inaccessible to RV surveys. Finally, they generally require that the system be monitored for at least one full period of the companion. Thus, although RV surveys have been ongoing for over a decade, they are only now becoming sensitive to Jupiter-analogs (Marcy et al. 2003).

¹See <http://cfa-www.harvard.edu/planets/catalog.html> for a catalog and discovery references

Microensing is an alternative method of detecting planetary companions that overcomes many of the difficulties inherent in the classical methods. Mao & Paczyński (1991) first proposed that microensing could be used to detect planets; their ideas were subsequently expanded on by Gould & Loeb (1992). Soon after these first two seminal theoretical papers, several microensing planet searches were initiated (Pratt et al. 1995, Albrow et al. 2000), and microensing searches have now been ongoing, in some form, for nearly a decade. Here I review the basic theoretical concepts behind planetary microensing, and describe how microensing planet searches work in practice. I then give an overview of what the analyses of actual datasets have taught us about the practicality of the method itself, and about planetary companions in general. Finally, I briefly speculate on future prospects for microensing planet searches.

2. Microensing and Planets

Toward the Galactic bulge, microensing occurs when a compact object, typically a low-mass star, passes close to our line-of-sight to a more distant star. The lens splits the source into two images separated by $\sim 2\theta_E$, where

$$\theta_E \equiv \sqrt{\frac{4GM}{Dc^2}} = 300\mu\text{as} \left(\frac{M}{0.3M_\odot} \right)^{1/2}, \quad (1)$$

is the Einstein ring radius of the lens, D is defined by, $D \equiv D_{os}D_{ol}/D_{ls}$, and D_{os} , D_{ol} , and D_{ls} are the distances between the observer-source, observer-lens, and lens-source, respectively. Thus for typical microlenses toward the bulge, the images are unresolved, and the only observable is the sum of the magnification of the two images of the source, which depends (only) on the angular separation between the lens and source in units of θ_E . Since observer, source, and lens are in relative motion, this angular separation, and thus the magnification, will be a function of time: a microensing event. Normal microensing events have a characteristic, symmetric, three-parameter form. The parameters are the time of maximum magnification, the impact parameter, and the characteristic timescale of the event,

$$t_E \equiv \frac{\theta_E}{\mu} = 20 \text{ days} \left(\frac{M}{0.3M_\odot} \right)^{1/2}, \quad (2)$$

where μ is the relative lens-source proper motion. The timescale is the only parameter that carries any information about the lens. Figure 1 illustrates the image geometry as a function of the angular separation of the source, and the resultant microensing lightcurve.

If the primary lens has a planetary companion with projected position near the path of one of the two images created in the primary event, the planet will deflect the light from the image, creating a perturbation atop the nominal single lens curve. See Figure 1. This perturbation is the signature of a planetary companion to the primary lens. Three parameters describe the magnitude and shape of the perturbation, and hence define the observables. These are the mass ratio q , the instantaneous projected separation d between the planet and star, and the angle α of the source trajectory with respect to the planet-star axis.




Figure 1. Left: The images (dotted ovals) are shown for several different positions of the source (solid circles), along with the primary lens (dot) and Einstein ring (long dashed circle). If the primary lens has a planet near the path of one of the images, i.e. within the short-dashed lines, then the planet will perturb the light from the source, creating a deviation to the single lens light curve. Right: The magnification as a function of time is shown for the case of a single lens (solid) and accompanying planet (dotted) located at the position of the X in the top panel. If the planet was located at the + instead, then there would be no detectable perturbation, and the resulting light curve would be identical to the solid curve.

The most interesting parameter is q , which is related to and derived from the duration of the perturbation, $t_p \simeq \sqrt{q}t_E$, which is $\simeq 1$ day for $M_p = M_J$. The orientation and phase of the planetary orbit, as well as θ_E , are all generally unknown; thus d gives only statistical information about the semi-major axis a . Since the planet must be near to one (or both) of the two images in order to be detected, and the images are near the Einstein ring radius while the source is significantly magnified, microlensing is most sensitive to planets with $d \sim 1$, which corresponds to $a \sim D_{ol}\theta_E \simeq 2\text{AU} (M/M_\odot)^{1/2}$. Finally, α is a randomly distributed, geometric parameter, and is of no physical interest.

For fixed q and d , only certain values of α will lead to detectable perturbations (see Fig. 1). Therefore integration over α defines a geometric detection probability. This detection probability is roughly $\sim A(\theta_p/\theta_E) \simeq 15\%(q/10^{-3})^{1/2}$, where A is the instantaneous magnification and $\theta_p \equiv \sqrt{q}\theta_E$. Thus the detection probability is substantial, and high-magnification events are intrinsically more sensitive to planetary companions (Griest & Safizadeh 1998, Rattenbury et al. 2003).

The advantages of microlensing are several. First, no flux is needed from either the lens or source, so systems with distances of many kiloparsecs can be probed. Second, it is sensitive to planets at separations $a = 1 - 10$ AU

immediately, without monitoring for the entire orbital period. Finally, since the duration of the signal drops only as $\sqrt{q}$, it is possible to extend the sensitivity to low masses. The primary disadvantages of microlensing are that follow-up of the detected systems will be difficult due to their large distances, the observations are non-repeatable, and the durations of the perturbations are short. These last two disadvantages combine to yield the requirement for successful detection of planets via microlensing: nearly continuous, densely sampled, and reasonably accurate photometry of the primary microlensing events.

3. Alerts and Follow-up

The probability that a star toward the Galactic bulge will be microlensed at any given time is of order (10^{-6}). Therefore, survey collaborations (EROS, Afonso et al. 2001; MOA, Bond et al. 2001; OGLE, Udalski et al. 2000) must monitor several millions of stars at a time in order to detect ongoing events. Practically, this means that several fields are monitored on a once-per-night basis, and the photometry is generally not optimized for any given microlensing event. Therefore, current survey photometry is generally not sufficient to detect planetary companions. However, real-time data reduction enables these collaborations to issue alerts, public notification of ongoing events.^{2,3,4} This allows follow-up collaborations (EXPORT, Tsapras et al. 2001; MPS, Rhie et al. 2000; PLANET, Albrow et al. 2000) to monitor only those events that are ongoing at any given time, and to optimize sampling rates and exposures times to individual events.

An example of this two-tier procedure is shown in Figure 2. On the left is the OGLE lightcurve for the alert OGLE-1998-BUL-14. On the right is the PLANET photometry of the same event (Albrow et al. 2000). While the OGLE dataset is relatively sparse, the median sampling interval for the PLANET data is about 1 hour, or $10^{-3}t_E$, with very few gaps greater than 1 day. The 1σ scatter in I over the peak of the event (where the sensitivity to planets is the highest) is 1.5%. The dense sampling and excellent photometry means that the efficiency to detect massive companions in this event should be quite high. In fact, a rigorous search for planetary companions, and quantification of the detection efficiency (Gaudi & Sackett 2000), reveals that, although the dataset is indeed quite sensitive to M_J planets at separations of $a = 1 - 5$ AU, there is no evidence for a companion. Analyses of similar datasets of other events using a variety of techniques have clearly demonstrated that the requisite photometric accuracy and sampling needed to robustly detect planetary perturbations is readily achievable (Albrow et al. 2000, Rhie et al. 2000, Bond et al. 2002, Tsapras et al. 2002). Therefore microlensing is not only sensitive to $\sim M_J$ companions between 1 – 10 AU in principle, but also in practice.

²For EROS alerts, see <http://www-dapnia.cea.fr/Phys/Spp/Experiences/EROS/alertes.html>.

³For MOA alerts, see <http://www.roe.ac.uk/~iab/alert/alert.html>.

⁴for OGLE alerts, see <http://www.astrouw.edu.pl/~ftp/ogle/ogle3/ews/ews.html>.




Figure 2. Left: The top panel shows the magnification as a function of time for OGLE data of microlensing event OGLE-1998-BUL-14. The dashed line indicates the best-fit point-lens model. The bottom panel shows the residuals from the best-fit model. Right: The top panel shows the PLANET data for the same event. The bottom panel shows the residuals (Albrow et al. 2000).

4. Limits on Galactic Planets

Analyses of datasets of individual microlensing events have demonstrated that microlensing is a practical method to detect $\sim M_J$ planetary companions, and in some cases have ruled out companions to individual lenses over a large range of separations. However, a coherent, uniform analysis of a significant sample of events is necessary to draw conclusions about the population of companions to lenses as a whole. The first such study was performed by the PLANET collaboration, who analyzed their complete database of photometric measurements of bulge microlensing events during the first five years of their campaign, from 1995 to 1999 (Gaudi et al. 2002). After applying cuts to eliminate nearly equal-mass binaries, poorly sampled events, and poorly-constrained events, a well-defined sample of 43 events was analyzed.

Each event was searched for planetary perturbations. There were no clear detections. For each event, the detection efficiency was computed for a broad range of q and d , using the method of Gaudi & Sackett (2000). These efficiencies were then combined to produce an upper limit to the fraction $f(d, q)$ of lenses with a companion. The 95% confidence level (c.l.) upper limit to $f(d, q)$ from the PLANET five year analysis is shown in Figure 3.

Although microlensing constraints are the most robust and least model-dependent in the (d, q) -plane, this parameterization makes it difficult to compare with other methods, which are primarily sensitive to M_p and a . To convert from (d, q) to (a, M_p) , it is necessary to specify the typical mass M and physical Einstein ring radius $r_E = D_{ol}\theta_E$ of the primaries, and also to marginalize over all possible orientations and orbital phases (as d is the instantaneous projected separation). It is likely that the majority of the microlensing primaries are M




Figure 3. The left panel shows contours of the 95% confidence upper limit to the fraction of lenses with a companion as a function of the mass ratio q and projected separation d of the companion from the PLANET 5-Year analysis (Albrow et al. 2001). The right panel is a census of planetary companions to solar-type stars, as well as sensitivities of various detection methods. The exclusion region from the PLANET microlensing survey is shown as the triangular hatched regions; $< 45\%$, $< 33\%$ (outer, inner) of M dwarfs in the bulge have companions in these regions.

dwarfs in the Galactic bulge (Gould 2000, Albrow et al. 2001). Thus $M \simeq 0.3M_\odot$ and $r_E \simeq 2\text{AU}$. With these values, $q = 0.001 \Rightarrow m_p = 0.3M_J$, and $d = 1 \Rightarrow r_p = 2\text{AU}$. Figure 3 shows the resulting constraints in the (a, M_p) -plane.

The PLANET analysis demonstrates that $< 33\%$ of M dwarfs in the Galactic bulge have Jupiter-mass companions with separations between 1.5 and 4 AU, and less than 45% have $3M_J$ companions between 1 and 7 AU. The range of parameter space constrained by PLANET observations partially overlaps that surveyed by RV studies; indeed many of the planets detected by RV surveys fall within the PLANET 45% exclusion region. However, the fractions of local stars with planets in this range is $\sim 5\%$, and so the RV results are not in conflict with microlensing constraints. Indeed, as RV and microlensing surveys are probing disjoint populations, both in regards to the typical mass of the primaries ($M \sim M_\odot$ for RV versus $M \sim 0.3M_\odot$ for microlensing), and the parent populations (local stars for RV versus bulge stars for microlensing), a comparison between RV and microlensing results would likely provide clues as to how planet formation and survival depends on the primary mass and local environment. As I discuss in the next section, although microlensing constraints have not yet reached the fractions currently probed by RV surveys, continued monitoring should allow a meaningful comparison between the two methods in the next few years.

5. Future Prospects

In the immediate future, microlensing searches for extrasolar planets are likely to continue with the current two-tier “survey/follow-up” system. How can such searches be improved, and the rate at which statistics are obtained increased? Some improvement can be achieved by more efficient monitoring of events and/or by employing image subtraction, rather than PSF-fitting, photometry. However, the most crucial factor in the efficiency of the survey/follow-up system is the number and quality of the alerts provided by the survey collaborations. A large number of alerts allows the follow-up collaborations to monitor only those events that are the most promising and most efficient for the detection of planetary perturbations, i.e. bright, high-magnification events. Previous to 2002, $\lesssim 100$ alerts per year were provided by the various survey collaborations. However, the OGLE collaboration recently upgraded its camera system (Udalski et al. 2002), and an extrapolation from the number of events alerted so far this year suggests that ~ 500 per year can be expected. Assuming photon-limited photometry, optimal observational strategies, and that a fraction $f(q)$ of lenses have companions with mass ratio q , and separations uniformly distributed in the range $d = 0.5 - 2.5$, the rate $\mathcal{R}_{det}$ at which planetary detections can be expected is roughly,

$$\mathcal{R}_{det} \sim 1 \text{ yr}^{-1} \left(\frac{q}{10^{-3}} \right)^{1/2} \left(\frac{\mathcal{R}_{alert}}{500 \text{ yr}^{-1}} \right)^{1/2} \left(\frac{f(q)}{5\%} \right). \quad (3)$$

Thus current survey/follow-up microlensing searches should probe companion frequencies of a few percent over the next several years.

Microlensing is unique among planet detection methods in that, to zeroth order, the amplitude of the signal does not decrease as q decreases. Therefore, very low-mass planets are in principle detectable via microlensing. The ultimate limit is set by the size of the source star: if the angular size of the source θ_* is larger than θ_p , the signal will be ‘washed out’ by the finite size of the source. For main-sequence sources, this limit is reached at about $q \simeq 10^{-5}$, the mass ratio between an Earth mass planet and the typical microlensing primary. The duration and detection probability both decrease as $\sqrt{q}$. For $q = 10^{-5}$, the duration of the perturbation is ~ 1 hr and the detection probability is a few percent (Bennett & Rhie 1996)

Even if the photometric accuracies and sampling rates required to detect the perturbations from Earth-mass planets are achieved, a naive extrapolation of equation (3) suggests that it will be quite difficult to build up reasonable statistics on Earth-mass planets using the survey/follow-up method. This is simply due to the low detection probability and the finite total exposure time. Ground-based detection of Earth-mass planets might be possible via a ‘next-generation’ ground-based microlensing planet search. Such a search dispenses with the survey/follow-up model. Instead, it employs a network of 2m-class telescopes distributed throughout the Southern hemisphere equipped with large-format CCDs that continually monitor a few fields toward the bulge at extremely high cadence to simultaneously detect *and* monitor events for planets.

The primary difficulties with ground-based surveys for Earth-mass planets are weather, which hampers the ability to robustly detect and constrain the planetary perturbations, and atmospheric seeing, which causes main-sequence

sources to be severely blended even in exceptional conditions, thereby increasing the photometric noise. A space-based mission would overcome these difficulties. The Galactic Exoplanet Survey Telescope (GEST), a proposed 2m aperture telescope with a ~ 2 square degree field-of-view would provide interesting constraints on Earth-mass planets, including Earth analogs (Bennett & Rhie 2002). The primary disadvantage of a space-based survey is the high cost.

Both next-generation ground-based surveys and space-based missions would also be sensitive to very wide separation and free-floating planets (Di Stefano & Scalzo 1999a,b). Microlensing is the only method yet proposed to detect low-mass free-floating planets.

Acknowledgments. I would like to thank the SOC and LOC for organizing an enjoyable and productive conference. Support for this work was provided by NASA through a Hubble Fellowship grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555.

References

- Afonso, C. et al. 2001, *A&A*, 378, 1014
 Albrow, M. et al. 2000, *ApJ*, 535, 176
 Albrow, M. et al. 2001, *ApJ*, 556, L113
 Bennett, D. P. & Rhie, S. H. 1996, *ApJ*, 472, 660
 Bennett, D. P. & Rhie, S. H. 2000, *ApJ*, 574, 985
 Bond, I. A. et al. 2001, *MNRAS*, 327, 868
 Bond, I. A. et al. 2002, *MNRAS*, 333, 71
 Di Stefano, R. & Scalzo, R. A. 1999a, *ApJ*, 512, 564
 Di Stefano, R. & Scalzo, R. A. 1999b, *ApJ*, 512, 579
 Gaudi, B. S. & Sackett, P. D. 2000, *ApJ*, 528, 56
 Gaudi, B. S. et al. 2002, *ApJ*, 566, 463
 Gould, A. 2000, *ApJ*, 535, 928
 Gould, A. & Loeb, A. 1992, *ApJ*, 396, 104
 Griest, K. & Safizadeh, N. 1998, *ApJ*, 500, 37
 Mao, S., & Paczyński, B. 1991, *ApJ*, 374, 37
 Marcy, G. W. et al. 2003, *ApJ*, submitted (astro-ph/0207294)
 Pratt et al. 1995, in *IAU Symp. 173: Astrophysical Applications of Gravitational Lensing*, eds. C. S. Kochanek & J. N. Hewitt (Dordrecht: Kluwer), 221
 Rattenbury et al. 2003, these proceedings
 Rhie, S. H. et al. 2000, *ApJ*, 533, 378
 Tsapras, Y. et al. 2001, *MNRAS*, 325, 1205
 Tsapras, Y. et al. 2002, *MNRAS*, in press (astro-ph/0207086)
 Udalski, A. et al. 2000, *Acta Astronomica*, 50, 1
 Udalski, A. et al. 2002, *Acta Astronomica*, 52, 1